

CATTLE VACCINE GUIDE						BACTERIAL DISEASES										VIRAL DISEASES														
	REG. NR.	PACK SIZE	DOSE	ADMINISTRATION	Botulism	Bovine Brucellosis (<i>Brucella abortus</i>)	Anthrax	Pasteurella infections (<i>Mannheimia haemolytica</i>)	Blackquarter/Quarter evil (<i>C. chauvoei</i>)	Bacterial red urine (<i>C. haemolyticum</i> / <i>C. novyi</i> type D)	Black disease (<i>C. novyi</i> type B)	Redgut (<i>C. perfringens</i> type A)	Redgut (<i>C. perfringens</i> type B)	Redgut (<i>C. perfringens</i> type C)	Redgut (<i>C. perfringens</i> type D)	Gas gangrene (<i>C. septicum</i>)	Gas gangrene (<i>C. sordelli</i>)	Tetanus (<i>C. tetani</i>)	<i>E.coli</i>	Infectious Keratoconjunctivitis (<i>Moraxella bovis</i>)	BVD (Bovine viral diarrhoea)	IBR (Infectious bovine rhinotracheitis)	Rotavirus	Coronavirus	Lumpy skin disease	Paratyphoid (<i>Salmonella dublin</i> / <i>Salmonella typhimurium</i>)	PI3			
SINGLE																														
Anthrax [®]	G2064	100 ml	1 ml	s/c			•																							
Botuvax [®]	G2193	100 ml	2 ml	s/c	•																									
Sponsvax [®]	G2063	100 ml	2 ml	s/c					•																					
Bovilis [®] S	G3763	20 ml/100 ml	2 ml	s/c																							•			
Bovi-Tect P	G3002	10 ml/100 ml	1 ml	s/c			•																							
Lumpyvax [®]	G3673	20 ml/100 ml	1 ml	s/c																					•					
RB-51	G3056	10 ml/50 ml	2 ml	s/c		•																								
Piliguard [®] Pinkeye - 1 Trivalent	G2803	20 ml/100 ml	2 ml	i/m s/c																•										
COMBINATIONS																														
Multiclos	G3392	100 ml/250 ml	5 ml	s/c					•	•	•		•	•	•	•	•	•												
Covexin [™] -10	G3354	100 ml	2 ml	s/c					•	•		•	•	•	•	•	•	•												
Botuthrax	G3783	100 ml	2 ml	s/c	•		•																							
Blanthrax [®]	G1593	150 ml	2 ml	s/c			•																							
Duovax	G2328	100 ml	2 ml	s/c	•				•																					
Supavax [®]	G2643	100 ml	2 ml	s/c	•		•																							
Rotavec [®] Corona	G2955	40 ml	2 ml	s/c															•				•	•						
Bovi-Tect PI	G3001	10 ml/100 ml	1 ml	s/c				•														•								
Bovi-Tect III	G3211	100 ml	1 ml	s/c			•															•	•							
Respiravax [™]	G3867	20 ml/100 ml	1 ml	s/c			•															•	•							

BEES ENTSTOFGIDS						BAKTERIËLE SIEKTES										VIRUSSIEKTES														
	REG. NR.	VERPAKKING	DOSIS	TOEDIENING	Lamsiekte	Besmetlike misgeboorte (<i>Brucella abortus</i>)	Miltsiekte	Pasteurella infeksies (<i>Mannheimia haemolytica</i>)	Sponssiekte (<i>C. chauvoei</i>)	Bakteriese rooi urine (<i>C. haemolyticum</i> / <i>C. novyi</i> tipe D)	Aansteeklike nekrotiese hepatitis (<i>C. novyi</i> tipe B)	Rooiderm (<i>C. perfringens</i> tipe A)	Rooiderm (<i>C. perfringens</i> tipe B)	Rooiderm (<i>C. perfringens</i> tipe C)	Rooiderm (<i>C. perfringens</i> tipe D)	Kwaadaardige edeem (<i>C. septicum</i>)	Kwaadaardige edeem (<i>C. sordelli</i>)	Klem-in-die-kaak (<i>C. tetani</i>)	<i>E.coli</i>	Besmetlike Keratoconjunctivitis (<i>Moraxella bovis</i>)	BVD (Bees virus diarrree)	IBR (Besmetlike bees rinotrageitis)	Rotavirus	Coronavirus	Knopvelsiekte	Paratifus (<i>Salmonella dublin</i> / <i>Salmonella typhimurium</i>)	PI3			
ENKEL																														
Anthrax [®]	G2064	100 ml	1 ml	o/h			•																							
Botuvax [®]	G2193	100 ml	2 ml	o/h	•																									
Sponsvax [®]	G2063	100 ml	2 ml	o/h					•																					
Bovilis [®] S	G3763	20 ml/100 ml	2 ml	o/h																							•			
Bovi-Tect P	G3002	10 ml/100 ml	1 ml	o/h			•																							
Lumpyvax [®]	G3673	20 ml/100 ml	1 ml	o/h																					•					
RB-51	G3056	10 ml/50 ml	2 ml	o/h		•																								
Piliguard [®] Pinkeye - 1 Trivalent	G2803	20 ml/100 ml	2 ml	b/s o/h																•										
KOMBINASIES																														
Multiclos	G3392	100 ml/250 ml	5 ml	o/h					•	•	•		•	•	•	•	•	•												
Covexin [™] -10	G3354	100 ml	2 ml	o/h					•	•		•	•	•	•	•	•	•												
Blanthrax [®]	G1593	150 ml	2 ml	o/h			•																							
Botuthrax	G3783	100 ml	2 ml	o/h	•		•																							
Duovax	G2328	100 ml	2 ml	o/h	•				•																					
Supavax [®]	G2643	100 ml	2 ml	o/h	•		•																							
Rotavec [®] Corona	G2955	40 ml	2 ml	b/s															•				•	•						
Bovi-Tect PI	G3001	10 ml/100 ml	1 ml	o/h				•														•								
Bovi-Tect III	G3211	100 ml	1 ml	o/h			•															•	•							
Respiravax [™]	G3867	20 ml/100 ml	1 ml	o/h			•															•	•							

SHEEP VACCINE GUIDE

BACTERIAL DEASESES

	PACK SIZE	DOSE	ADMINISTRATION	Blackquarter (<i>C. chauvoei</i>)	Bacterial red urine (<i>C. haemolyticum/C. novyi</i> type D)	Black Disease (<i>C. novyi</i> type B)	Bloodgut (<i>C. perfringens</i> type A)	Lam dysentery (<i>C. perfringens</i> type B)	Bloodgut (<i>C. perfringens</i> type C)	Pulpy Kidney (<i>C. perfringens</i> type D)	Uterine gangrene (<i>C. septicum</i>)	Gas gangrene (<i>C. sordelii</i>)	Enzootic abortion (<i>Chlamydia abortus</i>)	Tetanus (<i>C. tetani</i>)	Botulism	Anthrax (<i>B. anthracis</i>)	<i>Mannheimia haemolytica</i> (YRP Technology)	<i>Pasteurella trehalosi</i> (YRP Technology)
SINGLE																		
Anthravax®	100 ml	0.5 ml	s/c															
Botuvax®	100 ml	1 ml	s/c															
Sponsvax®	100 ml	1 ml	s/c	•														
Pulpyvax®	100 ml/100d 250 ml/250d	1 ml	s/c							•								
Pulpyvax® 1 Shot	100 ml/100d	1 ml	i/m							•								
Ovillis® Enzovax	40 ml/20d	2 ml	s/c or i/m										•					
COMBINATIONS																		
Multiclos	100 ml/250 ml	3 ml	s/c	•	•	•		•	•	•	•	•						
Blanthrax®	150 ml	2 ml	s/c	•												•		
Botuthrax	100 ml	2 ml	s/c												•	•		
Duovax	100 ml	2 ml	s/c	•											•	•		
Supavax®	100 ml	2 ml	s/c	•											•	•		
Multivax P	100/250/500 ml	2 ml	s/c	•						•	•			•			•	
Multivax P Plus	100/250/500 ml	2 ml	s/c	•		•		•	•	•	•			•			•	•
Covexin™-10	100 ml	1 ml	s/c	•	•	•	•	•	•	•	•	•	•	•				

SKAAP ENTSTOFGIDS

BAKTERIËLE SIEKTES

	VERPAKKING	DOSIS	TOEDIENING	Sponsiekte (<i>C. chauvoei</i>)	Bakteriese rooi urine (<i>C. haemolyticum/C. novyi</i> type D)	Aansteeklike nekrotiese hepatitis (<i>C. novyi</i> type B)	Rooierm (<i>C. perfringens</i> type A)	Bloedpens (<i>C. perfringens</i> type B)	Bloedderm (<i>C. perfringens</i> type C)	Bloednier (<i>C. perfringens</i> type D)	Baarmoedersponsiekte (<i>C. septicum</i>)	Kwaadaardige edeem (<i>C. sordelii</i>)	Ensoötiese Aborsie (<i>Chlamydia Abortus</i>)	Klem-in-die-kaak (<i>C. tetani</i>)	Lamsiekte (<i>C. botulinum</i>)	Miltsiekte (<i>B. anthracis</i>)	<i>Mannheimia haemolytica</i> (YRP Technologies)	<i>Pasteurella trehalosi</i> (YRP tegnologie)	
ENKEL																			
Anthravax®	100 ml	0.5 ml	o/h														•		
Botuvax®	100 ml	1 ml	o/h												•				
Sponsvax®	100 ml	1 ml	o/h	•															
Pulpyvax®	100 ml/100d 250 ml/250d	1 ml	o/h							•									
Pulpyvax® 1 Shot	100 ml/100d	1 ml	b/s							•									
Ovillis® Enzovax	40 ml/20d	2 ml	o/h of b/s										•						
KOMBINASIES																			
Multiclos	100 ml/250 ml	3 ml	o/h	•	•	•		•	•	•	•	•							
Blanthrax®	150 ml	2 ml	o/h	•												•			
Botuthrax	100 ml	2 ml	o/h												•	•			
Duovax	100 ml	2 ml	o/h	•											•	•			
Supavax®	100 ml	2 ml	o/h	•											•	•			
Multivax P	100/250/500 ml	2 ml	o/h	•						•	•			•			•		
Multivax P Plus	100/250/500 ml	2 ml	o/h	•		•		•	•	•	•			•			•	•	
Covexin™-10	100 ml	1 ml	o/h	•	•	•	•	•	•	•	•	•	•	•					

VACCINATION SCHEDULE FOR HORSES

FOR TECHNICAL ADVISORS

DISEASE	PROPOSED VACCINATIONS					
	FOALS/WEANLINGS	YEARLINGS	PERFORMANCE HORSES	PLEASURE HORSES	BROODMARES	VACCINES
TETANUS	1st dose: 3 - 4 months 2nd dose: 4 - 5 months	Annual	Annual	Annual	Annual: 4 - 6 weeks prefoaling	Equilis® Prequenza TE
INFLUENZA	1st dose: 4 - 6 months 2nd dose: 4 - 7 months 3rd dose: 9 - 11 months	Annual	Annual	Annual	Annual	Equilis® Prequenza Equilis® Prequenza TE Equilis® Resequin
LOCAL REGULATIONS MUST BE ADHERED TO						
RHINOPNEUMONITIS	1st dose: 3 - 4 months 2nd dose: 4 - 5 months 3rd dose: 6 - 11 months	Biannual	Biannual	Biannual	Biannual at least one booster 4 - 6 weeks before foaling	Equilis® Resequin
STRANGLES (<i>Streptococcus equi</i>)	1st dose: 4 months 2nd dose: 5 months	Biannual	3 - 6 months	3 - 6 months		Equilis® StrepE
RABIES	3 - 4 months	Annual	Annual	Annual	Annual	Nobivac® Rabies
BOTULISM	3 - 4 months	Annual	Annual	Annual	Annual	Botuvax®

PERDE INENTINGSPROGRAM

VIR TEGNIESE ADVISEURS

SIEKTE	VOORGESTELDE INENTINGSPROGRAM					
	VULLENS	JAARROUD	PRESTASIEPERDE	RYPERDE	TEELMERRIES	ENTSTOF
KLEM-IN-DIE-KAAK	1ste dosis: 3 - 4 maande 2de dosis: 4 - 5 maande	Jaarliks	Jaarliks	Jaarliks	Jaarliks: 4 - 6 weke voor vul	Equilis® Prequenza TE
PERDEGRIEP	1ste dosis: 4 - 6 maande 2de dosis: 4 - 7 maande 3de dosis: 9 - 11 maande	Jaarliks	Jaarliks	Jaarliks	Jaarliks	Equilis® Prequenza Equilis® Prequenza TE Equilis® Resequin
MOET OOK AAN PLAASLIKE REGULASIES VOLDOEN WORD						
RINOPNEUMONITIS	1ste dosis: 3 - 4 maande 2de dosis: 4 - 5 maande 3de dosis: 6 - 11 maande	Halfjaarliks	Halfjaarliks	Halfjaarliks	Halfjaarliks of ten minste 4 - 6 weke voor vul	Equilis® Resequin
NUWESIEKTE (<i>Streptococcus equi</i>)	1ste dosis: 4 maande	Halfjaarliks	3 - 6 maande	3 - 6 maande		Equilis® StrepE
HONDSOLHEID	3 - 4 maande	Jaarliks	Jaarliks	Jaarliks	Jaarliks	Nobivac® Rabies
LAMSIEKTE	3 - 4 maande	Jaarliks	Jaarliks	Jaarliks	Jaarliks	Botuvax®