

## FOR EXTERNAL ANIMAL USE ONLY

### ACTIVYL® SPOT-ON SOLUTION

Reg. no. G4021 (Act 36/1947)

Namibia Reg. No. V14/18.3.10/1248 NSO

### INDICATIONS

**Activyl® Spot-On Solution** is used in the treatment and prevention of flea infestation (*Ctenocephalides felis*). **Activyl® Spot-On Solution** can be used as part of a treatment strategy for flea allergy dermatitis (FAD). Developing stages of fleas (eggs and larvae) in the pet's immediate surroundings are killed following contact with **Activyl® Spot-On Solution** treated pets.

### TARGET SPECIES

Dogs and Cats

### COMPOSITION

Contains: Indoxacarb 195 mg/ml.

Each unit dose (pipette) of **Activyl® Spot-On Solution** delivers:

	Unit Dose (mℓ)	Indoxacarb (mg)
<b>Activyl® Spot-On Solution</b> for toy dogs	0,51	99,45
<b>Activyl® Spot-On Solution</b> for small dogs	0,77	150,15
<b>Activyl® Spot-On Solution</b> for medium dogs	1,54	300,30
<b>Activyl® Spot-On Solution</b> for large dogs	3,08	600,60
<b>Activyl® Spot-On Solution</b> for extra-large dogs	4,62	900,90
<b>Activyl® Spot-On Solution</b> for small cats	0,51	99,45
<b>Activyl® Spot-On Solution</b> for large cats	1,03	200,85

### STORAGE

- Store at or below 30 °C.
- Do not remove the pipette from the packaging until ready to use.
- Do not use after the expiry date stated on the foil.

### WARNINGS

- **Withdrawal period:** Not applicable.
- **Activyl® Spot-On Solution** is highly flammable. Keep away from heat, sparks, open flame or other sources of ignition.
- Do not eat, drink or smoke while handling the product.
- This product contains indoxacarb. People with known hypersensitivity to indoxacarb should not use this product. Avoid contents coming into contact with eyes. If this occurs, rinse slowly and gently with water. Wash hands immediately after use.
- The safety of **Activyl® Spot-On Solution** has not been established in puppies and kittens younger than 8 weeks of age, and in pregnant or lactating bitches and queens. The responsible veterinarian should make a benefit-risk assessment before using **Activyl® Spot-On Solution** in these cases.
- The safety of **Activyl® Spot-On Solution** has not been established in dogs weighing less than 1,5 kg and in cats weighing less than 0,6 kg.
- Care should be taken to apply the dose to an area where the animal cannot lick it off, to ensure that animals do not groom each other following treatment, and that the content of the pipette or the applied dose does not come into contact with the eyes of the recipient and/or other animals. The product remains effective following shampoo treatment, water immersion (swimming, bathing) and exposure to sunlight.
- Dispose of all empty containers as per local waste disposal regulations and do not reuse for any other purpose.

- KEEP OUT OF REACH OF CHILDREN, ANIMALS AND UNINFORMED PERSONS
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, notify the registration holder.

## CONTRA-INDICATIONS

None

## ADVERSE REACTIONS

A brief period of hyper salivation may occasionally occur in cats if the animal licks the application site immediately after treatment. This is not a sign of intoxication and disappears within some minutes without treatment. Correct application will minimize licking of the application site. In rare cases of licking the product, gastro-intestinal signs (e.g. emesis, anorexia) or reversible neurological signs (e.g. incoordination, tremor, ataxia and mydriasis) have been observed in cats. If you notice any serious effects or other effects not mentioned in this package insert, please inform your veterinary surgeon.

## DIRECTIONS FOR USE- USE ONLY AS DIRECTED

### DOSAGE

**Dogs:** The recommended dose is 15 mg indoxacarb/kg bodyweight, equivalent to 0,077 ml/kg bodyweight. The following table defines the size of pipette to be used according to the weight of the dog:

Weight of dog (kg)	Pipette size to be used	Volume (ml)	Indoxacarb (mg/kg)
≤ 6,5	<b>Activyl® Spot-On Solution</b> for toy dogs	0,51	Minimum of 15
6,6 to 10	<b>Activyl® Spot-On Solution</b> for small dogs	0,77	15 to 23
10,1 to 20	<b>Activyl® Spot-On Solution</b> for medium dogs	1,54	15 to 30
20,1 to 40	<b>Activyl® Spot-On Solution</b> for large dogs	3,08	15 to 30
40,1 to 60	<b>Activyl® Spot-On Solution</b> for extra-large dogs	4,62	15 to 23

For dogs weighing more than 60 kg, the appropriate combination of pipettes should be used.

**Cats:** The recommended dose is 25 mg indoxacarb/kg bodyweight, equivalent to 0,128 ml/kg bodyweight. The following table defines the size of pipette to be used according to the weight of the cat:

Weight of cat (kg)	Pipette size to be used	Volume (ml)	Indoxacarb (mg/kg)
≤ 4	<b>Activyl® Spot-On Solution</b> for small cats	0,51	Minimum of 25
> 4	<b>Activyl® Spot-On Solution</b> for large cats	1,03	Minimum of 50

Following a single administration, **Activyl® Spot-On Solution** will prevent further flea infestation for at least 4 weeks.

For optimal control of flea infestation, **Activyl® Spot-On Solution** can be safely administered at 2 to 4 week intervals throughout the flea season.

### Advice on Correct Administration

Open 1 foil pouch and remove the pipette.


Ensure that the animal is calm and standing still for easy application.

### Step 1: Opening of Pipette

With the 1 hand, hold the pipette in an upright position away from your face and with the other hand, snap tip open by bending it and folding it back on itself.



### Step 2: Application of Activyl® Spot-On Solution (Dogs)



With 1 hand, part the hair and place pipette tip against the skin between shoulder blades. Squeeze pipette firmly, applying the entire contents of the pipette(s) evenly to 3 or 4 spots, all located at different application sites along the dog's backline from the shoulder to the base of the tail. Do not apply an excessive amount of solution at any one spot that could cause some of the solution to run off the side of the animal.

### Step 2: Application of Activyl® Spot-On Solution (Cats)



With 1 hand, part the hair and place pipette tip against the skin at the base of the skull. Squeeze pipette firmly, applying the entire contents of the pipette(s).

### PRESENTATION

**Activyl® Spot-On Solution** is packaged in individual-dose pipettes in an outer foil package. It is supplied in units of 1, 4 or 6 pipettes per cardboard box.

#### REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.  
20 Spartan Road, Spartan  
1619, RSA  
Tel: +27 (0) 11 923 9300  
Fax: +27 (0) 11 392 3158  
[www.msd-animal-health.co.za](http://www.msd-animal-health.co.za)

#### MANUFACTURER

Merck Animal Health  
2667 West Dual Street, Baton Rouge  
Louisiana 70814-4906, USA

### DATE OF PUBLICATION OF PACKAGE INSERT

27 August 2015