

SUGGESTED BROILER VACCINATION PROGRAM IN NORMAL CHALLENGE AREAS

Day of Age	Disease	Vaccine	Application Route
Day 1 (Hatchery)	Infectious Bronchitis	Infectious Bronchitis vaccine	Coarse spray
	Newcastle Disease	NOBILIS® ND C2	Coarse spray
Day 10	Infectious Bronchitis Newcastle Disease	NOBILIS® IB MA5 + CLONE 30	Drinking water / coarse spray
Day 14	Gumboro Disease	NOBILIS® GUMBORO D78	Drinking water
Day 19	Gumboro Disease	NOBILIS® GUMBORO D78	Drinking water
Day 21	Newcastle Disease	NOBILIS® ND CLONE 30	Drinking water

If broilers are being kept longer than 42 days an additional Newcastle vaccine at 28 days of age is recommended.

SUGGESTED BROILER VACCINATION PROGRAM IN HIGH RISK GUMBORO AREAS

Day of Age	Disease	Vaccine	Application Route
Day 1 (Hatchery)	Infectious Bronchitis	Infectious Bronchitis vaccine	Coarse spray
	Newcastle Disease	NOBILIS® ND C2	Coarse spray
Day 10	Infectious Bronchitis Newcastle Disease	NOBILIS® IB MA5 + CLONE 30	Drinking water / coarse spray Drinking water / coarse spray
Day 11	Gumboro Disease	NOBILIS® GUMBORO 228E	Drinking water
Day 16	Gumboro Disease	NOBILIS® GUMBORO D78	Drinking water
Day 21	Newcastle Disease	NOBILIS® ND CLONE 30	Drinking water

If broilers are being kept longer than 42 days an additional Newcastle vaccine at 28 days of age is recommended.



BROILER VACCINE INFORMATION

NOBILIS® ND C2

REG. NO. G2872 (ACT 36/1947) | ZIMBABWE REG. NO. 2012/80.23.17/9680

NOBILIS® ND C2 is a live freeze-dried vaccine, indicated for the primary vaccination of fowls at one day of age against Newcastle Disease.



NOBILIS® MA5 + CLONE 30

REG. NO. G2301 (ACT 36/1947) | ZIMBABWE REG. NO. 94/80.23.10/9383

NOBILIS® MA5 + CLONE 30 is a live attenuated freeze-dried vaccine against Infectious Bronchitis and Newcastle Disease in fowls, and is used for the immunisation of healthy fowls against the Massachusetts type of serologically related types of Infectious Bronchitis and against Newcastle Disease. It is administered by the coarse spray or eye drop method from 1 day of age.



NOBILIS® ND CLONE 30

REG. NO. G2466 (ACT 36/1947) | ZIMBABWE REG. NO. 94/80.23.10/9363

NOBILIS® ND CLONE 30 is a live freeze-dried vaccine to prevent Newcastle Disease in fowls. It is administered by the coarse spray or eye drop method from 1 day of age.



NOBILIS® GUMBORO D78

REG. NO. G2483 (ACT 36/1947) | ZIMBABWE REG. NO. 94/80.23.10/9366

NOBILIS® GUMBORO D78 is a live freeze-dried vaccine for the immunisation of fowls against Infectious Bursal Disease (IBD / Gumboro). The vaccine can be administered to fowls from day 7 by drinking water.



NOBILIS® GUMBORO 228E

REG. NO. G2423 (ACT 36/1947) | ZIMBABWE REG. NO. 95/80.23.10/9389

NOBILIS® GUMBORO 228E is a live vaccine for the active immunisation of fowls against infectious Bursa Disease (IBD / Gumboro). The vaccine can be administered to fowls from day 7 by drinking water.



GENERAL GUIDELINES TO WATER VACCINATION

- Calculate the volume of water the birds will consume within 2 hours, use that quantity to reconstitute vaccine
- Thirst birds for 1 hr to 1 hr 45 mins before drinking water vaccination
- Stabilise the water to be used for vaccination, use Vac-safe dechlor tablets
- Open vaccine vial under the stabilised water
- Ensure vaccine is consumed within 2 hours
- Discard unconsumed water containing vaccine after 2 hours



Vac-Safe® dechlorinating tablets
Dechlorinating Agent (min) 1.25 g
FD&C Blue I Colourant (min) 1.5 g

Intervet South Africa (Pty) Ltd, Reg. No. 1991/006580/07
20 Spartan Road, Spartan, 1619, RSA | Private Bag X2026, Isando, 1600, RSA
Tel: +27 (0) 11 923 9300, **Fax:** +27 (0) 11 392 3158, **Sales Fax:** +27 (0) 86 603 1777
www.msd-animal-health.co.za | ZA/OPO/0715/0008

MSD
Animal Health