

FOR EXTERNAL ANIMAL USE ONLY

ECTODEX

Reg. No. G2965 (Act 36/1947)

Namibia Reg. No. V02/18.3.4/662 NS0

This vaccine may only be used by or under the supervision of persons authorised in terms of or authorised in terms of section 23 (1) (c) of the Veterinary and Para-Veterinary Professions Act, 1982 (Act No. 19/1982).

INDICATIONS

Controls demodectic mange, sarcoptic mange, ticks and lice.

COMPOSITION

Contains: Amitraz 12,5 % *m/v*

HARMFUL

STORAGE

- Store in a cool, dry place at or below 25 °C.

WARNINGS

- Do not use on Chihuahuas.
- Do not use on dogs suffering from heat stress.
- **Ectodex** is not recommended for use on cats.
- Transient drowsiness may be noted at the demodectic mange dose. Recovery is spontaneous.
- **KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.**
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- For use on dogs only.
- Wear rubber gloves when diluting and using the dip wash.
- Avoid contact with dip and wash immediately and thoroughly with soap and hot water after accidental contamination.
- Protect product from extreme heat and cold and from dampness.
- Tightly reseal the container if all the contents have not been used.
- Discard any unused dip wash and all empty containers according to local waste disposal regulations and do not reuse for any other purpose.

USE DURING PREGNANCY AND LACTATION

Teratology and reproduction studies in laboratory animals revealed no adverse effects.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED FOR EXTERNAL USE OF DOGS ONLY.

- **Ectodex** should not be mixed with any other dip.
- This dip wash must be freshly prepared prior to use.

Mixing instructions

Mix the required volume of **Ectodex** with the specified volume of lukewarm water in a plastic bucket.

	Ectodex	Water (lukewarm)
Demodectic mange	40 ml	10 l
Sarcoptic mange	20 ml	10 l
Ticks and lice	10 ml	10 l

Demodectic Mange: Repeat treatment 5 to 7 days later until no live mites can be identified, or continue treatment for at least 3 weeks after all clinical signs have subsided.

Sarcoptic Mange: Repeat treatment 7 to 10 days later.

Ticks and Lice: Repeat treatment as necessary.

PRESENTATION

100 ml bottles.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za

MANUFACTURER

Nulandis
14 Field Street
Lilianton, Boksburg

DATE OF PUBLICATION OF THIS PACKAGE INSERT

13 November 2001