

## FOR ANIMAL USE ONLY

4\*

### NASALCUR®

Reg. No. G1831 (Act 36/1947)

Namibia Reg. No. V93/18.1.3/519 NSO



### INDICATIONS

Remedy for nasal worm, liver fluke and wireworm in sheep and goats as well as liver fluke and wireworm in cattle.

### COMPOSITION

Contains: Rafoxanide 3,0 % *m/v*

\* Salicylanilide

### CAUTION

### STORAGE

Store in a cool dry place.

### WARNINGS

- **Withdrawal period:** Do not slaughter animals for human consumption within 28 days of dosing.
- Milk from treated animals must not be used for human consumption.
- Do not dose lactating animals whose milk or milk products are intended for human consumption.
- Care must be taken in calculating the correct dosage rate as overdoses may lead to toxicity.
- Always ensure that all drenching procedures are carefully carried out. Care should be exercised in the dosing and handling of animals.
- Dispose of empty containers according to local waste disposal regulations and do not use for any other purpose.
- KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

### PRECAUTIONS

No worm remedy, not even **Nasalcur®**, should be dosed to sheep and goats that have not been immunised against pulpy kidney disease (enterotoxaemia).

Do not treat animals more frequently than every 3 weeks.

### DIRECTIONS FOR USE - USE ONLY AS DIRECTED

Shake well before use.

Use as an oral drench.

### DOSAGE

**Sheep and goats -** Dose 2,5 ml per 10 kg live mass.

**Cattle -** Dose 12,5 ml per 50 kg live mass

**Example: Cattle, sheep and goats**

LIVEMASS	DOSE	NUMBER OF DOSES PER PACK			
		200 ml	1 l	5 l	10 l
10 kg	2,5 ml	80	400	2 000	4 000
20 kg	5 ml	40	200	1 000	2 000
30 kg	7,5 ml	27	133	666	1 333
40 kg	10 ml	20	100	500	1 000
50 kg	12,5 ml	16	80	400	800
100 kg	25 ml	8	40	200	400

## EFFICACY

WORM SPECIES	SHEEP AND GOATS		CATTLE	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Liver Fluke	6 weeks **	*		*
Giant Liver Fluke	8 weeks *	*		*

### Sheep and goats:

Nasal worm (*Oestrus ovis*): \* Controls all larval stages

### Explanation:

\* Control ( $\geq 90$  % effective)

\*\* Aids in control (60 to 89 % effective)

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

## PRESENTATION

200 ml, 1 l, 5 l, 10 l Plastic bottles

## REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.  
20 Spartan Road, Spartan  
1619, RSA  
Tel. +27 (0) 11 923 9300  
Fax. +27 (0) 11 392 3158  
[www.msd-animal-health.co.za](http://www.msd-animal-health.co.za)

## MANUFACTURER

Nulandis  
14 Field Street  
Lillianton, Boksburg

## DATE OF PUBLICATION OF THIS PACKAGE INSERT

27 September 2000