


FOR ANIMAL USE ONLY



NEM-A-RID® 3,75 %

Reg. No. G3548 (Act 36/1947)

Namibia Reg No. V05/18.1.8/789 NS0



INDICATIONS

Roundworm and liver fluke remedy for cattle, sheep and goats as well as a nasal worm remedy for sheep and goats.

Targets: wireworm, brown stomach worm, bankrupt worm, cattle bankrupt worm, long-necked bankrupt worm, large-mouthed bowel worm, hookworm, nodular worm, lungworm, liver fluke, giant liver fluke.

COMPOSITION

Contains: Levamisole Hydrochloride 3,75 % *m/v*

Rafoxanide 3,75 % *m/v*

* Imidazothiazole + Salicylanilide

CAUTION

STORAGE

Store in a cool, dry place.

WARNINGS

- **Withdrawal period:** Do not slaughter animals for human consumption within 28 days of dosing.
- Do not dose lactating animals whose milk or milk products are intended for human consumption.
- No worm remedy, not even **Nem-A-Rid® 3,75%** should be dosed to sheep and goats that have not been immunised against pulpy kidney (enterotoxemia).
- Care must be taken in calculating the correct dosage rate as overdose may lead to toxicity.
- Always ensure that all drenching procedures are carefully carried out. Care should be exercised in the dosing and handling of animals.
- Do not treat animals under the age of 6 weeks.
- Do not treat animals more frequently than every 3 weeks.
- In case of poisoning call a doctor and make this label available to him.
- Dispose of any empty containers according to local waste disposal regulations and do not reuse for any other purpose.
- **KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.**
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED

Shake well before use.

DOSAGE

Sheep and goats: 2 ml/10 kg live mass

Cattle: 10 ml/50 kg live mass

Example:

CATTLE, SHEEP AND GOATS

Live Mass	Dose	Number of doses per pack			
		200 ml	1 l	5 l	10 l
10 kg	2 ml	100	500	2 500	5 000
20 kg	4 ml	50	250	1 250	2 500
30 kg	6 ml	33	166	833	1 666
40 kg	8 ml	25	125	625	1 250
50 kg	10 ml	20	100	500	1 000
100 kg	20 ml	10	50	250	500

EFFICACY

Worm Species	Sheep and Goats		Cattle	
	Immature	Adult	Immature	Adult
Wireworm	*	*	*	*
Brown stomach worm	**	*		*
Bankrupt worm	*	*		
Cattle bankrupt worm			*	*
Long-necked bankrupt worm	*	*		
Large-mouthed bowel worm	*	*		
Hookworm	*	*		*
Nodular worm	*	*	**	*
Lungworm	**	**		
Liver fluke	6 weeks **	*		*
Giant liver fluke	8 weeks *	*		*

* Controls (≥ 90 % effective)

** Aids in the control of (60 to 89 % effective)

SHEEP AND GOATS

Nasal worm (*Oestrus ovis*) – Controls (> 90 % effective) all larval stages.

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

PRESENTATION

200 ml, 1 l, 5 l and 10 l plastic bottles.

REGISTRATION HOLDER

Intervet South Africa (Pty) Ltd.
20 Spartan Road, Spartan
1619, RSA
Tel: +27 (0) 11 923 9300
Fax: +27 (0) 11 392 3158
www.msd-animal-health.co.za

MANUFACTURER

Burchem Research (Pty) Ltd.
54 Electron Avenue
Isando

DATE OF PUBLICATION OF PACKAGE INSERT

14 October 2004