



GUIDELINES FOR RAISING CALVES IN FEEDLOTS

Scan QR codes for more product information



Arrival



Rest



Rehydration



Roughage

Rest phase

Rest phase must be for 24-36 hours

Immunovite

Minerals and Vitamins (35 ml / 100 l water 3-5 days)



24 hours later

Minerals

Covexin® 10

Clostridial diseases such as Blackquarter and Redgut or *Histophilus somni* and *Clostridia* spp

Bovilis® C7 Somni

Botuthrax®

Botulism/Anthrax

Bovilis® Vista Once

Lungs (BVD Tipe 1 & 2, IBR, PI3, BRSV, *Pasteurella multocida*, *Mannheimia haemolytica*)

Lumpyvax®

Lumpy Skin Disease

Revalor® XR and Revalor® H

(Revalor® XR starts working 70 days after implantation)

Reverin LA 230 (summer)

For the treatment of tick-borne gallsickness, heartwater, respiratory disease, footrot, and navel-ill

Procaine benzylpenicillin 150 mg/ml, Benzathine benzylpenicillin 125 mg/ml (winter)

Consult a veterinarian

180 mg Tildipirosin (winter)

Consult a veterinarian

Sovereign®

Deworming / Liver fluke and ticks or

Gardal 10 %

Deworming / Liver fluke and Tapeworm

Taktic® Cattle Spray or Delete® All

External parasites

10 - 14

Day 10 - 14

Covexin® 10

Must be followed up again after 6 weeks (42 days) with **Covexin® 10**

or

Bovilis® C7 Somni

Should be followed up after 3-4 weeks with **Bovilis® C7 Somni**

Botuvax®

Must be administered 4-6 weeks after vaccination with **Botuthrax®**



Zilmax®

Last 30-35 days for completion (72 hour withdrawal)



Lumpyvax® G3673



Immunovite V24724



Revalor® XR G4490



Bovilis® C7 Somni G3114



Taktic® Cattle Spray G2535



Revalor® H G2023



Reverin LA G3521



Botuthrax® G3783



Sovereign® G3831



Zilmax® G2180



Covexin® 10 G3354



Bovilis® Vista Once SQ G4061



Gardal 10 % G3201



Botuvax® G2193



Delete® All G2837