



GUIDELINES FOR RAISING CALVES IN FEEDLOTS

Scan QR
codes for
more product
information



Arrival



Rest



Rehydration



Roughage

Rest phase

Rest phase must be for 24-36 hours

Immunovite

Minerals and Vitamins (35 ml / 100 l water 3-5 days)



24 hours later

Minerals

Covexin® 10

Clostridial diseases such as Blackquarter and Redgut **or** *Histophilus somni* and *Clostridia* spp

Bovilis® C7 Somni

Botuthrax®

Botulism/Anthrax

Bovilis® Vista Once

Lungs (BVD Tipe 1 & 2, IBR, PI3, BRSV, *Pasteurella multocida*, *Mannheimia haemolytica*)

Lumpyvax®

Lumpy Skin Disease

Revalor® XR and

Revalor® H

(Revalor® XR starts working 70 days after implantation)

Reverin LA 230

(summer)

For the treatment of tick-borne gallsickness, heartwater, respiratory disease, footrot, and navel-ill

Procaine benzylpenicillin

Consult a veterinarian

150 mg/ml, Benzathine

benzylpenicillin

125 mg/ml (winter)

180 mg Tildipirosin

(winter)

Consult a veterinarian

Sovereign®

Deworming / Liver fluke and ticks **or**

Gardal 10 %

Deworming / Liver fluke and Tapeworm

Taktic® Cattle Spray

External parasites

or Delete® All

10-14

Day 10 - 14

Covexin® 10

Must be followed up again after 6 weeks (42 days) with **Covexin® 10**

or

Bovilis® C7 Somni

Should be followed up after 3-4 weeks with **Bovilis® C7 Somni**

Botuvax®

Must be administered 4-6 weeks after vaccination with **Botuthrax®**



Zilmax®

Last 30-35 days for completion (72 hour withdrawal)



Lumpyvax®
G3673



Immunovite
V24724



Revalor® XR
G4490



Bovilis® C7
Somni
G3114



Taktic®
Cattle Spray
G2535



Revalor® H
G2023



Reverin LA
G3521



Botuthrax®
G3783



Sovereign®
G3831



Zilmax®
G2180



Covexin® 10
G3354



Bovilis® Vista
Once SQ
G4061



Gardal 10 %
G3201



Botuvax®
G2193



Delete® All
G2837