



GUIDELINES FOR RAISING CALVES ON THE FARM

Scan QR
codes for
more product
information

Field Preparation



Arrival



Rest



Rehydration



Roughage



Day 0

Minerals

Vit A

Covexin® 10

Bovilis® C7 Somni

Lumpyvax®

Botuthrax®

Bovilis® Vista Once SQ

Ralgro® Cattle or

Revalor® XS

Reverin LA 230

(summer)

Procaine benzylpenicillin

150 mg/ml, Benzathine

benzylpenicillin

125 mg/ml (winter)

Sovereign®

Gardal 10 %

Fluxacur NF

Tactic® Cattle Spray

or Delete® All

Blackquarter/Malignant Oedema/Clostridium **or**
Histophilus somni and Clostridium

Lumpy Skin Disease

Botulism/Anthrax

Lungs (BVD Tipe 1 & 2, IBR, PI3, BRSV, *Pasteurella*
multocida, *Mannheimia haemolytica*)

90 days on field (to grow skeletal frame)

200 days

For the treatment of tick-borne gallsickness,
heartwater, respiratory disease, footrot, and navel-ill

Consult with veterinarian

Deworming / Liver fluke and ticks **or**

Deworming / Liver fluke and tapeworm **or**

Deworming / Liver fluke

External parasites



Day 18 - 30

Covexin® 10

or

Bovilis® C7 Somni

Botuvax®

Must be followed up with
Covexin® 10 after 6 weeks (42 days).

Must be followed up with
Bovilis® C7 Somni after
3-4 weeks.

Must be administered 4-6
weeks after **Botuthrax®**.

Zilmax®

Last 30-35 days with
72 hour withdrawal



Lumpyvax®
G3673



Ralgro® Cattle
G1406



Bovilis® C7
Somni
G3114



Revalor® XS
G3925



Tactic®
Cattle Spray
G2535



Revalor® H
G2023



Reverin
LA 230
G3521



Botuthrax®
G3783



Sovereign®
G3831



Zilmax®
G2180



Covexin® 10
G3354



Bovilis® Vista
Once SQ
G4061



Gardal 10 %
G3201



Botuvax®
G2193



Delete® All
G2837



Fluxacur NF
G3202